Common <u>Dragonflies</u> and <u>Damselflies</u> of the Chicago Region

Volunteer Stewardship Network – Chicago Wilderness

Produced by: John & Jane Balaban, Jennie Kluse & Robin Foster, with assistance of Laurel Ross and support from the Gordon & Betty Moore Foundation. Photos © John & Jane Balaban; [balx2@comcast.net] North Branch Restoration Project, with additions by © Thomas Murray (27, 32) and © Vincent Hickey (30). © Environmental & Conservation Programs, The Field Museum, Chicago, IL 60605 USA. [http://www.fmnh.org/chicagoguides/]. Chicago Wilderness Guide #1 version 2 (4/2006)

LIBELLULIDAE - Skimmers

Large, showy, frequently seen resting on or flying low over vegetation. Often hunt from a perch like Kingbirds. Also includes our smallest dragonflies (Nannothemis and Perithemis) and the ubiquitous Meadowhawks.



1 Sympetrum rubicundulum ♀/♂

Ruby Meadowhawk: male and female mating in "wheel" position.

RESOURCES:

Drangonflies of Indiana by J. R. Curry. Indiana Academy of Science. 2001.

ISBN: 1-883362-11-3

Beginner's Guide to Dragonflies by Nikula and Sones with D. and L. Stokes. Little, Brown, and Company.

2002. ISBN: 0-316-81679-5

Damselflies of the Northeast by E. Lam. Biodiversity

Books, 2004, ISBN: 0-9754015-0-5

Damselflies of the North Woods by B. DuBois. Kollath-Stensaas Pub. 2005. ISBN: 0-9673793-7-7

http://bugguide.ent.iastate.edu/node/view/191/bgimage http://cirrusimage.com/dragonflies.htm http://wisconsinbutterflies.org/damselflies/



2 Sympetrum obtrusum ♂



3 Sympetrum vicinum



4 Sympetrum semicinctum ♂

White-faced Meadowhawk: white face. 32-36mm Yellow-legged Meadowhawk: yellow legs. 30-36mm Band-winged Meadowhawk: half amber wings. 26-38mm Above species are medium-sized and common. Adult males are usually red; females and immature males are yellow; both sexes sometimes with black abdominal markings.



5 Nannothemis bella

Elfin Skimmer: our tiniest dragonfly; female has wasp-like pattern on abdomen



6 Perithemis tenera

Eastern Amberwing: male has amber wings.



7 Perithemis tenera

Eastern Amberwing: female has spotted wings.

19-25mm



8 Celithemis elisa

Calico Pennant: female has same pattern but golden colors



Celithemis eponina

Halloween Pennant: sits on tips of vegetation even on Four-spotted Skimmer: stout body; amber and black 28-34mm the windiest days.



10 Libellula quadrimaculata ♂

34-39mm wing markings. 42-49mm

Common **Dragonflies** and **Damselflies** of the Chicago Region

Volunteer Stewardship Network – Chicago Wilderness

Produced by: John & Jane Balaban, Jennie Kluse & Robin Foster, with assistance of Laurel Ross and support from the Gordon & Betty Moore Foundation. Photos © John & Jane Balaban; [balx2@comcast.net] North Branch Restoration Project, with additions by © Thomas Murray (27, 32) and © Vincent Hickey (30). © Environmental & Conservation Programs, The Field Museum, Chicago, IL 60605 USA. [http://www.fmnh.org/chicagoguides/]. Chicago Wilderness Guide #1 version 2 (4/2006)



11 Leucorrhinia intacta ♂

Dot-tailed Whiteface: bright white face; yellow dot on Wandering Glider: never seems to land; flies with Anax Blue Dasher: forward swept wings; white face; top of male abdomen.



14 Erythemis simplicicollis

Tramea onusta

similar T. carolina.

on abdomen.

Red-mantled Saddlebags: red wing patches have clear

ď

Common Whitetail: male/female dimorphism in wing

area near body; this clear area is larger than in the



12 Pantala flavescens

27-36mm swarm; clear wings with orange stigma.



15 Erythemis simplicicollis ♂

Eastern Pondhawk: female is green with dark markings Eastern Pondhawk: males turn rich blue with age.



18 Tramea lacerata

Black Saddlebags: large black patches at base of hindwings only.



46-55mm

21 Plathemis lydia

each wing; yellow slash markings on abdomen.



13 Pachydiplax longipennis ♂

45-51mm seagreen eyes; yellow and brown striped thorax. 32-45mm



16 Libellula vibrans

Great Blue Skimmer: long wings with black tips; 39-48mm blue-green eyes; white face. 50-63mm



19 Libellula luctuosa

Widow Skimmer: black basal patches on all wings; males also have white patch in middle of wings.

36-49mm



22 Libellula pulchella ♂

Common Whitetail: females have three brown spots in Twelve Spotted Skimmer: three black spots in each wing; solid yellow stripe on each side of abdomen.

pattern; males have broad black stripes in center of wing. 39-47mm 39-47mm 38-54mm

Common <u>Dragonflies</u> and <u>Damselflies</u> of the Chicago Region

Volunteer Stewardship Network – Chicago Wilderness

Produced by: John & Jane Balaban, Jennie Kluse & Robin Foster, with assistance of Laurel Ross and support from the Gordon & Betty Moore Foundation. Photos © John & Jane Balaban; [balx2@comcast.net] North Branch Restoration Project, with additions by © Thomas Murray (27, 32) and © Vincent Hickey (30). © Environmental & Conservation Programs, The Field Museum, Chicago, IL 60605 USA. [http://www.fmnh.org/chicagoguides/]. Chicago Wilderness Guide #1 version 2 (4/2006)

AESHNIDAE - Darners

Large bodied, with brilliant blue and green colors, unmarked wings. Strong non-stop flyers, eat on the wing like swallows. \Rightarrow

MACROMIIDAE - Cruisers

Large, black and yellow markings, green eyes, long legs, low rapid flight. ↓



23 Aeshna canadensis 🗸



24 Anax junius

Canada Darner: genus generally brown striped with blue Common Green Darner: recently emerged male; mature on thorax; the many species makes ID difficult. 62-73mm male is bright blue; CBS "eye" on head.



Macromia illinoiensis

Swift River Cruiser: newly emerged.



26 Epitheca cynosura &

Common Baskettail: early flight season; brown with yellow markings; wings clear except for dark patch at 65-75mm base 38-44mm



27 Epitheca princeps

Prince Baskettail: largest emerald; often misidentified as Libellula pulchella but rarely lands. ©Thomas Murray



28 Dorocordulia libera 🕜

Racket-tailed Emerald: small dark dragonfly with brilliant green eyes; abdomen spoon-shaped.



29 Somatochlora hineana

Hine's Emerald: the many Striped Emeralds makes species ID difficult; this one is federally endangered.



Slender brown dragonflies often with metallic green or bronze sheen. Some species have bright green eyes which can be seen in flight.



Usually perch horizontally. Recognized by widely spaced eyes. Abdomen often inflated at tip. Many very similar species in the family. ↓



30 Dromogomphus spinosus



31 Arigomphus submedianus o'



32 Stylurus plagiatus o

Black-shouldered Spinyleg: long spines mixed with Jade Clubtail: moderately wide reddish-brown abdomen Russet-tipped Clubtail: reddish-brown club; blue eyes; shorter ones on long hind leg. ©Vincent J Hickey 55-65mm tip; male claspers yellow; green eyes. 51-55mm long slender body. ©Thomas Murray 55-63mm

Common <u>Dragonflies</u> and <u>Damselflies</u> of the Chicago Region

Volunteer Stewardship Network – Chicago Wilderness

Produced by: John & Jane Balaban, Jennie Kluse & Robin Foster, with assistance of Laurel Ross and support from the Gordon & Betty Moore Foundation. Photos © John & Jane Balaban; [balx2@comcast.net] North Branch Restoration Project, with additions by © Thomas Murray (27, 32) and © Vincent Hickey (30). © Environmental & Conservation Programs, The Field Museum, Chicago, IL 60605 USA. [http://www.fmnh.org/chicagoguides/]. Chicago Wilderness Guide #1 version 2 (4/2006)



Dragonflies spend most of their lives in the water as larvae.



34 Eclose process

The animal emerges from the larval skin after metamorphosis as a soft-skinned "teneral" adult.



35 Exuviae

The larval skin left behind when the new adult flies away.



Males and females dimorphic. Can often identify females by identifying a nearby male of the same species. There are many damselflies in the region, but these eight images allow ID to family, and perhaps to genus. \Rightarrow



36 Calopteryx maculata ♀/♂

a butterfly; no stigma; sits horizontally on leaves. Male and female in "wheel" position 39-57 mm



37 Archilestes grandis &

Ebony Jewelwing: large size; broad wings; flutters like Great Spreadwing: very large; long stigma; wings held wide, dragonfly-like; hangs vertically on stems.

50-62mm

38 Lestes dryas ♂

Emerald Spreadwing: emerald green; wings held open but swept back; long stigma; hangs vertically on stems. 32-40mm



39 Lestes rectangularis

Slender Spreadwing: male with long abdomen relative to wings; vein pale at wing tip.



40 Argia apicalis

Blue-fronted Dancer: small stigma; wings folded over back but held above the body; lands horizontally; can be 37-53mm found dancing among sunny spots in trails.



41 Enallagma civile ♂

Familiar Bluet: many species, mostly blue-black; wings held down alongside body; "helicopters" up and down along stems. 28-39mm



42 Enallagma cyathigerumり/♂

Northern Bluet: male and female mating in "wheel" position.



43 Ischnura verticalis o

Eastern Forktail: very small, jewel-like; flies among low vegetation like Bluets.

29-40mm 21-29mm